

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/266742622>

26. Valdeyernos, Toledo Mountains (central Spain)

Article in Grana · October 2014

DOI: 10.1080/00173134.2014.936490

CITATIONS

4

READS

78

3 authors, including:



Miriam Dorado-Valiño

44 PUBLICATIONS 758 CITATIONS

[SEE PROFILE](#)



José Antonio López-Sáez

Spanish National Research Council

534 PUBLICATIONS 7,424 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



The Hinojos Project [View project](#)



The prehistoric ceremony center of Castillejo del Bonete: one of the most important archaeological discoveries of the 21st century // El centro ceremonial prehistórico de Castillejo del Bonete: uno de los descubrimientos arqueológicos más importantes del siglo XXI. [View project](#)

PROOF COVER SHEET

Journal acronym: SGRA

Author(s): Miriam Dorado-Valiño, José Antonio López-Sáez and Enrique García-Gómez

Article title: 26. Valdeyernos, Toledo Mountains (central Spain)

Article no: 936490

Enclosures: 1) Query sheet
2) Article proofs

Dear Author,

1. Please check these proofs carefully. It is the responsibility of the corresponding author to check these and approve or amend them. A second proof is not normally provided. Taylor & Francis cannot be held responsible for uncorrected errors, even if introduced during the production process. Once your corrections have been added to the article, it will be considered ready for publication.

Please limit changes at this stage to the correction of errors. You should not make trivial changes, improve prose style, add new material, or delete existing material at this stage. You may be charged if your corrections are excessive (we would not expect corrections to exceed 30 changes).

For detailed guidance on how to check your proofs, please paste this address into a new browser window:
<http://journalauthors.tandf.co.uk/production/checkingproofs.asp>

Your PDF proof file has been enabled so that you can comment on the proof directly using Adobe Acrobat. If you wish to do this, please save the file to your hard disk first. For further information on marking corrections using Acrobat, please paste this address into a new browser window:
<http://journalauthors.tandf.co.uk/production/acrobat.asp>

2. Please review the table of contributors below and confirm that the first and last names are structured correctly and that the authors are listed in the correct order of contribution. This check is to ensure that your name will appear correctly online and when the article is indexed.

Sequence	Prefix	Given name(s)	Surname	Suffix
1		Miriam	Dorado-Valiño	
2		José Antonio	López-Sáez	
3		Enrique	García-Gómez	

Queries are marked in the margins of the proofs, and you can also click the hyperlinks below.

AUTHOR QUERIES

General points:

1. **Permissions:** You have warranted that you have secured the necessary written permission from the appropriate copyright owner for the reproduction of any text, illustration, or other material in your article. Please see <http://journalauthors.tandf.co.uk/permissions/usingThirdPartyMaterial.asp>.
2. **Third-party content:** If there is third-party content in your article, please check that the rightsholder details for re-use are shown correctly.
3. **Affiliation:** The corresponding author is responsible for ensuring that address and email details are correct for all the co-authors. Affiliations given in the article should be the affiliation at the time the research was conducted. Please see <http://journalauthors.tandf.co.uk/preparation/writing.asp>.
4. **Funding:** Was your research for this article funded by a funding agency? If so, please insert 'This work was supported by <insert the name of the funding agency in full>', followed by the grant number in square brackets '[grant number xxxx]'.
5. **Supplemental data and underlying research materials:** Do you wish to include the location of the underlying research materials (e.g. data, samples or models) for your article? If so, please insert this sentence before the reference section: 'The underlying research materials for this article can be accessed at <full link> / description of location [author to complete]'. If your article includes supplemental data, the link will also be provided in this paragraph. See <http://journalauthors.tandf.co.uk/preparation/multimedia.asp> for further explanation of supplemental data and underlying research materials.
6. The **CrossRef database** (www.crossref.org/) has been used to validate the references. Mismatches will have resulted in a query.

AQ1 Please provide missing page numbers for the "Dorado-Valiño et al., 2014" references list entry.

AQ2 Please confirm NPPs as non-pollen palynomorphs, abbreviated in figure.

How to make corrections to your proofs using Adobe Acrobat/Reader

Taylor & Francis offers you a choice of options to help you make corrections to your proofs. Your PDF proof file has been enabled so that you can edit the proof directly using Adobe Acrobat/Reader. This is the simplest and best way for you to ensure that your corrections will be incorporated. If you wish to do this, please follow these instructions:

1. Save the file to your hard disk.
2. Check which version of Adobe Acrobat/Reader you have on your computer. You can do this by clicking on the “Help” tab, and then “About”.

If Adobe Reader is not installed, you can get the latest version free from <http://get.adobe.com/reader/>.

3. If you have Adobe Acrobat/Reader 10 or a later version, click on the “Comment” link at the right-hand side to view the Comments pane.

4. You can then select any text and mark it up for deletion or replacement, or insert new text as needed. Please note that these will clearly be displayed in the Comments pane and secondary annotation is not needed to draw attention to your corrections. If you need to include new sections of text, it is also possible to add a comment to the proofs. To do this, use the Sticky Note tool in the task bar. Please also see our FAQs here: <http://journalauthors.tandf.co.uk/production/index.asp>.

5. Make sure that you save the file when you close the document before uploading it to CATS using the “Upload File” button on the online correction form. If you have more than one file, please zip them together and then upload the zip file.

If you prefer, you can make your corrections using the CATS online correction form.

Troubleshooting

Acrobat help: <http://helpx.adobe.com/acrobat.html>

Reader help: <http://helpx.adobe.com/reader.html>

Please note that full user guides for earlier versions of these programs are available from the Adobe Help pages by clicking on the link “Previous versions” under the “Help and tutorials” heading from the relevant link above. Commenting functionality is available from Adobe Reader 8.0 onwards and from Adobe Acrobat 7.0 onwards.

Firefox users: Firefox’s inbuilt PDF Viewer is set to the default; please see the following for instructions on how to use this and download the PDF to your hard drive:

http://support.mozilla.org/en-US/kb/view-pdf-files-firefox-without-downloading-them#w_using-a-pdf-reader-plugin

CONTRIBUTIONS TO THE EUROPEAN POLLEN DATABASE

26. Valdeyernos, Toledo Mountains (central Spain)

MIRIAM DORADO-VALIÑO¹, JOSÉ ANTONIO LÓPEZ-SÁEZ¹ & ENRIQUE GARCÍA-GÓMEZ²

5 ¹Center of Human and Social Sciences, Spanish Council for Scientific Research, Madrid, Spain, ²Servicio de Medio Ambiente, Diputación de Toledo, Toledo, Spain

Site details

10 Valdeyernos bog (39° 26' 28" N, 4° 05' 47" W; 850 m above sea level (a.s.l.)) lies on the southern slope of the 'Sierra de Tornos' in the central part of the Toledo Mountains (Quintos de Mora Farm, Guadalerzas district, Los Yébenes, Toledo). The peat-bog extends over an area of 3.93 ha and it is a 'special protected habitat' in accordance with Law 9/1999 of Castilla-La Mancha on Nature Conservation. It was designated an 'environmental micro-reserve' in 2003 by the Decree 319/2003 (Martín-Herrero et al. 2004). The area experiences a Mediterranean climate with dry and warm summers, influenced by Atlantic winds from the southwest. The average annual temperature is 14–15 °C and the annual precipitation is 600–700 mm. The precipitation is irregular over the year with maximum values during winter (Perea & Perea 2008). The vegetation of the area is typically 25 Mediterranean, a mixed forest of deciduous oaks *Quercus pyrenaica* Willd. and *Q. faginea* subsp. *broteroi* (Cout.) A.Camus, accompanied by *Acer monspessulanum* L., *Arbutus unedo* L., *Q. ilex* subsp. *ballota* (Desf.) Samp., *Sorbus torminalis* (L.), *Crataegus monogyna* Jacq., *Phillyrea angustifolia* L., *Erica arborea* L., *E. scoparia* L. and *Cistus populifolius* L. (Gómez-Manzaneque 1988; Baonza et al. 2010). The bog vegetation is composed mainly of *Sphagnum palustre* L., *Drosera rotundifolia* L., *Pinguicula lusitanica* L., *E. tetralix* L., *Calluna vulgaris* (L.) Hull and *Molinia caerulea* (L.) Moench (Velasco 1980; Martín-Herrero et al. 2004). The bed-rock is an old siliceous basement made up mainly of sandstones, lutites and quartzites of Early Ordovician age (Muñoz-Jiménez 1976).

Sediment description

The profile is 50 cm deep and was collected from the central part of the bog in June 2006 by the last two authors using a Russian corer 5 cm in diameter. The composition was as follows:

0–14 cm: dark brown, partly decomposed *Sphagnum*-Cyperaceae peat with sand and gravels > 1 mm (Th2 Gmin3 Gmaj+ Dh1–10 YR 2/1) 45
14–34 cm: dark brown, decomposed *Sphagnum*-Cyperaceae peat with herbaceous and rather numerous wood fragments (Th2 Dh2 D11–10 YR 2/1) 50
34–50 cm: grey to olive silt with sand and gravels > 1 mm and macro-charcoal fragments (34–36, 39–43, 46–48 cm) (Ld1 Dh1 Dg1 D11 Gmaj2–5 YR 3/2)

Dating

Accelerator mass spectrometry (AMS) carbon-14 (¹⁴C) dating was performed on bulk peat samples by the 'Centro Nacional de Aceleradores' (CNA, CSIC) in Sevilla (Spain). The calibration (2σ range) was performed using CALIB 6.0. and CALIBomb (with the calibration dataset NH zone 1) software (Stuiver & Reimer 1993; Hua & Barbetti 2004). Calibrated radiocarbon dates are given as cal BC or cal AD. Dating results are as follows:

14–15 cm: CNA 717, pM 101.27 ± 0.26 (cal AD 1953–1956) 65
29–30 cm: CNA 719, 95 ± 30 BP (cal AD 1683–1930)
40 cm: CNA 718, 515 ± 30 BP (cal AD 1326–1444)
48–50 cm: CNA 093, 2935 ± 33 BP (1261–1027 cal BC)




Figure 1. Pollen diagram from Valdeyernos bog (Toledo Mountains, central Spain). Selected pollen, spores of ferns and non-pollen palynomorphs (NPPs) taxa are represented. Hollow curves are exaggerated 2×.

Percent Modern (pM) is used to describe radiocarbon measurements for the past 45 years (<http://www.radiocarbon.com/PDF/Beta-AMS-Methodology.pdf>)

Interpretation

The core was sampled at 2 cm intervals. More than 500 terrestrial pollen grains were identified in each sample. The pollen sum (100%) includes all pollen grains except for hygrophytes and aquatic plants, spores of ferns and non-pollen palynomorphs (Figure 1). The arboreal vegetation comprised *Betula*, *Corylus* and both deciduous and evergreen *Quercus* throughout the entire sequence. This vegetation was quite different from the close by Patateros bog (Dorado-Valiño et al. 2014), where the arboreal vegetation is scarce during the same time period. The differences between the two pollen records could be due to fact that the Patateros bog is located on a northern slope and the Valdeyernos bog on a southern slope. Therefore, the Valdeyernos site was more affected by south-western Atlantic warm and humid winds than the Patateros site; the Valdeyernos bog is, furthermore, located in a little valley where drain streams lead to high edaphic humidity. Four pollen assemblage zones were defined using CONISS in TILIA® and TILIA-GRAPH® (Eric C. Grimm) computer programs:

VD-I (50–40 cm; c. 1300 cal BC–cal AD 1420)

The landscape is dominated by a shrub community composed of *Erica*-type accompanied by *Cistus ladanifer*-type, *Helianthemum*-type, *Calluna vulgaris* and Poaceae. The high values of *Erica*-type pollen could be an overrepresentation of local species as the bog is large. This vegetation also contained isolated stands of *Betula*, *Corylus* and both deciduous and evergreen *Quercus* with an undergrowth of ferns (*Pteridium aquilinum* (L.) Kuhn, *Hymenophyllum*). The presence of hygrophytes (Cyperaceae), freshwater algae (Chrysophyceae, *Spirogyra*) and ascospores of HdV-18 and *Byssothecium circinans* Fuckel suggests moist conditions and relatively high water tables (Van Geel & Aptroot 2006). The maximum percentage of HdV-18 (48 cm) can be correlated with the onset of the so-called 2.8 ka cal BP event (c. 850–760 cal BC; Van Geel et al. 1998), while minimum values (46–44 cm, c. 230 cal BC–cal AD 275) with an arid interval described in the Iberian Peninsula at this time (Martin-Puertas et al. 2008).

VD-II (40–34 cm; c. cal AD 1420–1665)

In this pollen zone, the previous landscape progressively changed to a vegetation dominated by *Betula*, *Corylus* and deciduous *Quercus*, although heathers are still abundant. The chronological boundaries of this zone

correlate well with the onset of the Little Ice Age, which, in Spain, was a cold interval with an initial dry phase, until cal AD 1550, and a later more humid one until the recent present (Manrique & Fernández-Cancio 2000). Towards the end of the pollen zone, a maximum of *Chaetomium* is detected coinciding with a level of macro-charcoals and the appearance of *Olea europaea* L.

VD-III (34–10 cm; c. cal AD 1665–1970)

Groups of birch and hazel were distributed along the periphery of the bog, while the slopes of the surrounding hills were covered by mixed oak forests (probably *Quercus pyrenaica* and *Q. faginea*). *Olea europaea* and evergreen *Quercus* are scarcely represented. The presence of *Pinus pinaster* Aiton pollen during the last century (the uppermost 28 cm) was contributed by planted trees (Gómez-Manzaneque 1988). The non-arboreal vegetation is mainly composed of *Erica*-type and Poaceae with a similar representation of *Calluna vulgaris* and *Cistus ladanifer*-type. This pollen zone is characterized by maximum values of dung-related fungal spores (*Sordaria*, *Sporormiella*, *Cercophora*, *Podospora*, *Tripterospora*, *Coniochaeta*), although the values of anthropozoogenous taxa (*Chenopodiaceae*, *Plantago lanceolata*-type, *Plantago major*/*Plantago media*-type, *Urtica dioica*-type) are relatively low, suggesting local grazing (López-Sáez & López-Merino 2007). Weak human impact is suggested by the continuous presence of anthropogenic pollen indicators (*Cardueae*, *Cichorioideae*, *Rumex acetosella*-type). Erosive processes are reflected by the presence of *Glomus*, *Pseudoschizaea circula* (Wolff) Christopher and *Entorrhiza*. The progressive decline of *Betula*, hygrophytes, freshwater algae and *Byssothecium circinans* suggests initial drier conditions, which could favour the peat-bog formation at 34 cm, as shown by the sedimentary description and the continuous presence of *Tilletia sphagni* Navashin. Two maximum peaks for HdV-18 indicate wet phases (Mighall et al. 2006), while minimum values at 34 cm (cal AD 1665) and 28–24 cm (cal AD 1873–1895) can be correlated with two severe drought episodes reconstructed from roagation ceremonies recorded at the Toledo Cathedral (Domínguez-Castro et al. 2008).

VD-IV (10–2 cm; c. cal AD 1970–2006)

The landscape is almost the same as in the previous pollen zone but with a greater representation of *Erica*-type and a general decrease of coprophilous fungi.

Acknowledgements

This contribution was financed by Project CGL2011-30512-C02-01 of the R&D National

Plan of the Spanish Government, by the Technology Research ‘Consolider’ Programme (TCP-CSD2007-00058), and by the Specific Agreement Diputación de Toledo-CSIC.

References

- Baonza J, Caparrós R, García N, Martínez F, Gómez-Manzaneque F. 2010. Flora vascular de los Quintos de Mora (Los Yébenes, Toledo). *Ecología* 23: 39–58.
- Domínguez-Castro F, Santisteban JI, Barriendos M, Mediavilla R. 2008. Reconstruction of drought episodes for central Spain from roagation ceremonies recorded at the Toledo Cathedral from 1506 to 1900: A methodological approach. *Global and Planetary Change* 63: 230–242. doi:10.1016/j.gloplacha.2008.06.002.
- Dorado-Valiño M, López-Sáez JA, García-Gómez E. 2014. Contributions to the European Pollen Database 21: Patateros, Toledo Mountains (central Spain). *Grana* 53: XX–XX.
- Gómez-Manzaneque F. 1988. La cubierta vegetal en los Montes de Mora (Los Yébenes, Toledo). *Ecología* 2: 111–130.
- Hua Q, Barbetti M. 2004. Review of tropospheric bomb ¹⁴C data for carbon cycle modelling and age calibration purposes. *Radiocarbon* 46: 1273–1298.
- López-Sáez JA, López-Merino L. 2007. Coprophilous fungi as a source of information of anthropic activities during the Prehistory in the Amblés Valley (Ávila, Spain): The archaeopalynological record. *Revista Española de Micropaleontología* 39: 103–116.
- Manrique E, Fernández-Cancio A. 2000. Extreme climatic events in dendroclimatic reconstructions from Spain. *Climatic Change* 44: 123–138.
- Martín-Herrero J, Frontaura MC, Plaza F. 2004. Microrreservas de Castilla-La Mancha. *Medio Ambiente en Castilla-La Mancha* 11: 1–48.
- Martín-Puertas C, Valero-Garcés BL, Mata MP, González-Samperiz P, Bao R, Moreno A, Stefanova V. 2008. Arid and humid phases in southern Spain during the last 4000 years: The Zoñar lake record, Córdoba. *The Holocene* 18: 907–921. doi:10.1177/0959683608093533.
- Mighall TM, Martínez-Cortizas A, Biester H, Turner SE. 2006. Proxy climate and vegetation changes during the last five millennia in NW Iberia: Pollen and non-pollen palynomorph data from two ombrotrophic peat bogs in the North Western Iberian Peninsula. *Review of Palaeobotany and Palynology* 141: 203–223. doi:10.1016/j.revpalbo.2006.03.013.
- Muñoz-Jiménez J. 1976. Los Montes de Toledo. Estudio de Geografía Física. Oviedo: Instituto J.S. Elcano, CSIC and Universidad de Oviedo.
- Perea D, Perea R. 2008. Vegetación y flora de los Montes de Toledo. *Guía de Campo*. Toledo: Covarrubias.
- Stuiver M, Reimer PJ. 1993. Extended ¹⁴C database and revised CALIB radiocarbon calibration program. *Radiocarbon* 35: 215–230.
- Van Geel B, Aptroot A. 2006. Fossil ascomycetes in Quaternary deposits. *Nova Hedwigia* 82: 313–329. doi:10.1127/0029-5035/2006/0082-0313.
- Van Geel B, Van der Plicht J, Kilian MR, Klaver ER, Kouwenberg JHM, Renssen H, Reynaud-Farrera L, Waterbolk HT. 1998. The sharp rise of ¹⁴C ca. 800 cal BC: Possible causes, related climatic teleconnections and the impact on human environments. *Radiocarbon* 40: 535–550.
- Velasco A. 1980. Notas sobre los enclaves higroturbosos de los Montes de Toledo (España). *Anales del Jardín Botánico de Madrid* 37: 125–128.